



Osteo SAP



Science-based Osteo-minerals; vitamins and amino acids in non-GMO vegetable capsules

Osteoporosis is a multifactorial disease with roots in genetics, endocrine function, exercise and nutritional habits. Osteo' means bone and 'porosis', thinning or becoming more porous, together literally meaning 'thinning of bone'. Adequate nutrition is critical in osteoporosis prevention and treatment. Many nutritional factors have been examined for associations with osteoporosis and bone mass. Calcium is essential for bone development and stability and low calcium intake is believed to contribute to the development of osteoporosis. Vitamin D is essential for the maintenance of calcium levels and increases its uptake in the intestine. Boron is involved in calcium, magnesium and phosphorus metabolism contributing to prevention of bone loss associated with osteoporosis. Vitamin K1 from a vegetable source activates osteocalcin; the major non-collagen protein in bone and a specific marker of bone formation. Copper, manganese and zinc make up an essential trio in bone metabolism as cofactors for specific enzymes. Calcium, magnesium, phosphorus, trace minerals, and protein are components of bone tissue and are all found in microcrystalline hydroxyapatite complex (MCHC) in their natural ratios.

ACTIVE INGREDIENTS

Six vegetable capsules contain:

Microcrystalline Hydroxyapatite Complex (MCHC)	3500	mg
*Calcium (elemental from 3500mg MCHC)	875	mg
Phosphorus (naturally present in MCHC)	437	mg
Protein (naturally present in MCHC)	875	mg
*Magnesium oxide (elemental from 515 mg)	300	mg
*Magnesium bis-glycinate (elemental from 1500 mg)	150	mg
*Zinc picolinate	10	mg
*Manganese citrate	3	mg
Vitamin C ascorbic acid	50	mg
Vitamin K phylloquinone	100	mcg
Vitamin D3 cholecalciferol	1000	IU
*Boron citrate	3	mg
Horsetail 7% silica	25	mg
L-Taurine	50	mg
Glutamic acid HCL	50	mg
*Copper gluconate	1	mg

In non-GMO, water-based, vegetable capsules.

*ELEMENTAL DOSAGE

Osteo SAP contains **875mg of elemental calcium** and **450mg of elemental magnesium** a 2:1 ratio intended to provide bone support for individuals with some risk factors for bone loss.

Contains no: dairy products, preservative, artificial flavor or color.

Osteo SAP contains **180 vegetable capsules per bottle**.

ADULT DOSAGE

2 capsules with meals three times daily, or as directed by your health practitioner.

INDICATION

Independent use of **Osteo SAP** three times daily is expected to contribute to the uptake and maintenance of calcium and magnesium for the prevention of osteoporosis.

FEATURES

- **Osteo SAP** supplies a compatible and complementary blend of New Zealand bovine microcrystalline hydroxyl appetite (MCHC) calcium, minerals, vitamins and amino acids to enhance bone building osteoblasts, cells from which bones are formed.
- **Osteo SAP** contains the ideal elemental 2:1 ratio of calcium and magnesium with other cofactors to enhance calcium uptake for the prevention of osteoporosis.
- **Osteo SAP** supplies the most readily absorbable form of calcium phosphate (MCHC), the form found in human bones clinically proven to restore bone loss. It is free from pesticides, hormones, antibiotics and animal feed.

PURITY and CLEANLINESS

- **Osteo SAP** is third party tested to ensure freedom from heavy metals, PCB's, pesticides, volatile organics and impurities.



*Scientific Advisory Panel
(SAP):
adding nutraceutical
research to achieve
optimum health*

What is Osteoporosis?

Osteoporosis is a disease characterized by low bone mass, structural deterioration, diminished bone strength, and increased risk of fracture^{1,2}. Osteoporosis can be specifically related to bone mineral density (BMD) as there is a continuous inverse association between BMD and risk of fracture. In older adulthood, BMD and the risk for osteoporosis is characterized by the peak bone mass attained in adolescence and the rate of bone loss in later years. On average, women have lower BMD than men. Furthermore, heredity, age, ethnicity and a large number of environmental factors influence BMD and risk of osteoporotic fracture. In North America, 1 in 2 women and 1 in 4 men aged over 50 will fracture at some point in their life time². Currently, 1.5 million osteoporotic fractures of the hip, spine, wrist, and other sites occur each year, predominantly in postmenopausal white women. Moreover, hip and spine fractures often play a severe role in disability, dependence and increased risk of death¹.

Nutrition Therapy in the Prevention of Osteoporosis

Calcium – In adults, calcium accounts for 1 to 2 percent of human body weight with over 99 percent of total calcium found in bones and teeth. In bone, calcium is mainly present in the form of hydroxyapatite ($\text{Ca}_{10}(\text{PO}_4)_6(\text{OH})_2$), with calcium making up 40% of bone mineral and bone mineral contributing to 40% of bone weight^{1,3}. Bone is a dynamic tissue that is constantly undergoing osteoclastic bone resorption and osteoblastic bone formation. Calcium is necessary for bone health, is an important determinant of peak bone mass and thus helps prevent osteoporosis^{1,2}. Prospective studies indicate that increasing dietary calcium intake from 900 to 1500 mg daily may reduce the rate of bone loss in premenopausal women. Furthermore, controlled trials of supplementation with ≥ 500 mg of calcium (ideally in addition to 700 IU of vitamin D) daily for over a year in men and postmenopausal women demonstrate a reduction in rates of bone loss from the spine, hip, and total body. Discontinuation of supplemental calcium and vitamin D has been associated with a loss of BMD. Currently, Health Canada and the FDA have approved a health claim for dietary calcium and vitamin D, and a reduced risk of osteoporosis.

Phosphorus – Phosphorus is an essential mineral involved in bone mineralization, bone structure and most metabolic processes in the body¹. Approximately 80% of phosphorus in the body is found in bones and teeth as calcium phosphate crystals or hydroxyapatite. Furthermore, although calcium makes up 40% of bone mineral, phosphate accounts for nearly 60%. Thus phosphorus is fully as important for bone building as is calcium. Phosphorus is crucial for calcium absorption and utilization. Imbalance of the phosphorus-calcium ratio can interfere with calcium absorption, contributing to osteoporosis risk. Dietary phosphorus deficiency can limit osteoblast-mediated bone formation and enhance osteoclast-mediated bone resorption^{1,2}.

Protein – Protein is critical for a healthy skeleton and dietary protein plays an important role in bone health and the prevention of osteoporosis. A human study found that protein supplementation for 6 months in patients with recent hip fractures significantly lowered the rate of bone loss from the hip over a 1-year period¹. Additionally, the protective effect of dietary protein on bone loss seems to show most benefit in individuals with higher calcium intakes¹. The Framingham Osteoporosis Study revealed that elderly women and men with relatively lower protein intake had increased bone loss, supporting the importance of dietary protein in bone health maintenance in the elderly⁴.

Magnesium – Approximately half of the body's store of magnesium is found in bone tissue¹. Magnesium influences bone matrix and mineral metabolism, and magnesium depletion leads to cessation of bone growth, decreased osteoblastic and osteoclastic activity, osteopenia, and bone fragility⁵. Magnesium prevents bone fragility by directly affecting hydroxyapatite crystal formation and stabilizing amorphous calcium phosphate. Furthermore, magnesium regulates active calcium transport and metabolism. Magnesium deficiency can be a risk factor for postmenopausal osteoporosis³. In a 2-year study of menopausal women, magnesium therapy prevented fractures and led to a significant increase in bone density⁵.

Manganese, Zinc, and Copper – These 3 trace minerals are cofactors for enzymes essential to bone tissue synthesis and maintenance^{1,6}. During a 2-year trial of post-menopausal women, the effect of these 3 trace minerals on bone loss at the spine, alone or in combination with calcium was studied and revealed no decline in spine BMD as compared to a 3.5% loss of BMD in the control group¹.

Vitamin C – The formation or cross-linking of matrix proteins and collagen fibrils, necessary to build and maintain bones, cartilage and joints, is dependent on vitamin C as well as copper, zinc, and manganese¹. Interference with cross-linking results in structurally weak bone. Thus,

vitamin C is essential for proper calcification of bones (the process giving strength and hardness to bones). Epidemiological studies have observed a positive association between vitamin C intakes and bone mass, where low intakes of vitamin C was associated with increased rate of BMD loss. Similarly, another study revealed that higher vitamin C was inversely associated with fractures².

Vitamin K – Vitamin K is a fat-soluble vitamin that is involved in blood coagulation, is required for bone metabolism and assists in reducing urinary calcium excretion^{1,2}. Vitamin K1 (phylloquinone) is present in green leafy vegetables and other plants, whereas vitamin K2 (menaquinone) is present in liver, meats, and foods prepared by fermentation, such as cheeses^{1,2}. Vitamin K is a cofactor in the formation of the gamma-carboxylation of glutamyl residues in the bone proteins osteocalcin, matrix gamma-carboxyglutamyl protein, and protein S. Deficiency of each of these vitamin K-dependent proteins is associated with osteopenia^{1,6}. Studies have found that a high concentration of undercarboxylated osteocalcin and low dietary phylloquinone intake, resulting in low concentration of serum vitamin K, are associated with lower BMD and increased risk of hip fracture^{1,2}.

Vitamin D3 – Vitamin D is obtained from the diet in the forms of ergocalciferol (vitamin D2) and cholecalciferol (vitamin D3) as well as from cutaneous synthesis after exposure to sunlight^{1,2}. Vitamin D3 is modified in the liver and kidney to make 1,25-Dihydroxycholecalciferol (calcitriol), the most active metabolite of Vitamin D. The function of Vitamin D is to maintain serum calcium and phosphorus concentrations through regulating calcium absorption from the intestine or calcium resorption from bone, and vitamin D is necessary for maintenance of healthy bone. The role of vitamin D insufficiency in osteoporosis is strongly recognized. For men and women over 50 years of age, evidence suggests that the plasma level of 25(OH)D or calcidiol (clinical indicator of vitamin D status) needed to minimize fracture risk is ≥ 80 nmol/L and that an intake of 800-1000 IU/day of vitamin D3 is needed to bring the population average to this level¹. A higher intake of vitamin D2 would be needed to reach the same level.

Boron – Boron is a trace element that affects calcium, phosphorus and magnesium absorption and metabolism¹. An increase in boron intake results in a decrease in urinary excretion of calcium, phosphorus, and magnesium as well as a simultaneous increase in serum estradiol levels. Although the mechanism is unknown, it is proposed that boron is necessary for formation of certain steroid hormones or hydroxylation of 25(OH)D. Thus, dietary boron deficiency may lead to an increase in osteoporosis risk. Furthermore, research suggests a positive effect on bone health with 3 mg/day of boron supplementation².

Silicon – Silicon has been shown to be important in the formation and strengthening of bone and horsetail is a rich source of silicon. With dietary silicon depletion, both extracellular matrix (collagen) and bone mineral (hydroxyapatite) are compromised⁷. The average daily intake of silicon in the Western-world is ~ 20 -50mg/day and the major sources of dietary silicon are cereals/grains and some fruits and vegetables. In a recent epidemiological study, silicon intake positively correlated with BMD at 4 hip sites in men and premenopausal women but not in postmenopausal women. The study concluded that higher silicon intake (>40 mg/day) in men and younger women may positively affect skeletal health, particularly cortical bone health. Preliminary human study suggests benefit of silicon intake, but further research is needed before specific recommendations can be made.

Glutamic Acid and L-taurine – Hydrochloric acid (HCl) is produced in the stomach and is needed in ionization and absorption of some forms of calcium. Numerous studies observe a reduction in HCl gastric secretion with advancing age, which can be associated with a reduction in available calcium for absorption⁸. Glutamic acid is a source of HCl to assist in the breakdown and uptake of calcium. L-taurine is a non-essential amino sulfonic acid that also aids in the absorption of minerals, such as magnesium.

References

1. Shils ME. *Modern Nutrition in Health and Disease*. Tenth Edition 2006.
2. Nieves JW. *Am J Clin Nutr* 2005; 81:1232S-9S.
3. Institute of Medicine. 1997. DRIs for Calcium, Phosphorus, Magnesium, Vitamin D, and Fluoride. National Academy Press, Washington, D.C. USA.
4. Hannan MT, Tucker KL, Dawson-Hughes B, Cupples LA, et al. *J Bone Miner Res* 2000; 15:2504-12.
5. Sojka JE, Weaver CM. *Nutr Rev* 1995; 53(3):71-4
6. Institute of medicine. 2001. DRIs for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc. National Academy Press, Washington, D.C. USA.
7. Jugdaohsingh, R. et al. *J Bone Min Res* 2004; 19:297-307.
8. Kelly GS. *Alt Med Rev* 1997; 2(2):116-127.

Available from

[Dr. Ingrid Pincott, ND](#)

300-1170 Shopper's Row • Campbell River, BC • V9W 2C8 • Ph. (250) 286-3655 • (800) 898-6699 • Fax (250) 850-2078